

AD-A207 133

OFFICE OF NAVAL RESEARCH

Contract N0001489WX24082

R&T Code 4131011---04

Technical Report No. 11

EFFICIENT LASER ACTION FROM 1,3,5,7,8-PENTAMETHYLPYRROMETHENE-
BF₂ COMPLEX

by

Theodore G. Pavlopoulos, Mayur Shah, and Joseph H. Boyer

Prepared for Publication

in the

Optics Communications

NAVAL OCEAN SYSTEMS CENTER
San Diego, CA 92152

19 April 1989

Reproduction in whole or in part is permitted for any purpose of the United
States Government

This document has been approved for public release
and sale; its distribution is unlimited.

DTIC
ELECTE
APR 26 1989
S H D

089 4 25 189

ADA207133

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT This document has been approved for public release and sale; its distribution is unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) ONR-TR-11		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research, Chem. Div.	
6a. NAME OF PERFORMING ORGANIZATION Naval Ocean Systems Center		7b. ADDRESS (City, State and ZIP Code) Code 1113PS Arlington, VA 22217	
6b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N0001489WX24082	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		10. SOURCE OF FUNDING NUMBERS	
8b. ADDRESS (City, State and ZIP Code) Chemistry Division Code 1113PS Arlington, VA 22217		PROGRAM ELEMENT NO. PROJECT NO. TASK NO. AGENCY ACCESSION NO.	
11. TITLE (Include Security Classification) EFFICIENT LASER ACTION FROM 1,3,5,7,8-PENTAMETHYLPYRROMETHENE-BF ₂ COMPLEX			
12. PERSONAL AUTHOR(S) Theodore G. Pavlopoulos, Mayur Shah, and Joseph H. Boyer			
13a. TYPE OF REPORT Interim Technical		13b. TIME COVERED FROM 9/88 TO 9/89	
14. DATE OF REPORT (Year, Month, Day) 19 April 1989		15. PAGE COUNT	
16. SUPPLEMENTARY NOTATION To be published in Optics Communications			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD GROUP SUB-GROUP		Laser dyes, dye lasers, Pyrromethene-BF ₂ complexes, Boron difluoride, methanes, methyl 1,6-dicyano, pyrroles.	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) We observed efficient laser action under flashlamp excitation at 546 nm from 1,3,5,7,8-pentamethylpyrromethene-BF ₂ complex (PMP-BF ₂), dissolved in ethanol. This new laser dye lased about 300% more efficiently than Coumarin 545. The fluorescence, absorption (S-S), and triplet-triplet absorption spectra of PMP-BF ₂ were also recorded. To achieve greater water solubility the disodium 2,6-disulfonate derivative (PMPDS-BF ₂) was prepared. PMPDS-BF ₂ lases about 3/4 as efficiently as PMP-BF ₂ when dissolved in water and showed some photoinstability; however, in methanol it was highly resistant to photodegradation.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. T. G. Pavlopoulos		22b. TELEPHONE (Include Area Code) (619) 553-2792	
22c. OFFICE SYMBOL			

EFFICIENT LASER ACTION FROM 1,3,5,7,8-PENTAMETHYLPYRROMETHENE-BF₂ COMPLEX AND ITS DISODIUM 2,6,-DISULFONATE DERIVATIVE

Theodore G. Pavlopoulos, Marine Sciences and Technology Department
U.S. Naval Ocean Systems Center, San Diego, CA 92152


Mayur Shah and Joseph H. Boyer, Department of Chemistry
University of New Orleans, New Orleans, LA 70148

We observed efficient laser action under flashlamp excitation at 546 nm from 1,3,5,7,8-pentamethylpyrromethene-BF₂ complex (PMP-BF₂), dissolved in ethanol. This new laser dye lased about 300% more efficiently than coumarin 545. The fluorescence, absorption (S-S) and triplet-triplet absorption spectra of PMP-BF₂ were also recorded. To achieve greater water solubility the disodium 2,6-disulfonate derivative (PMPDS-BF₂) was prepared. PMPDS-BF₂ lases about 3/4 as efficiently as PMP-BF₂ when dissolved in water and showed some photo-instability; however, in methanol it was highly resistant to photodegradation.


1. INTRODUCTION

Recently we reported that flashlamp excitation of 1,3,5,7-tetramethylpyrromethene-BF₂ complex (TMP-BF₂) (2×10^{-4} M) in methanol brought about lasing emission (broad band BB) at 533 nm with about 10% greater efficiency than was obtained from coumarin 540A in ethanol (BB at 539 nm) [1]. Significantly, TMP-BF₂ had only a small amount of triplet-triplet (T-T) absorption over its fluorescence (laser action) spectral region. We speculated that even greater efficiency in laser action would be observed from pyrromethene complexes that show (1) extinction coefficients and quantum fluorescence yields

greater than those for TMP-BF₂ [ϵ_{505} (C₂H₅OH) = 83,200 and $Q_F = 0.80$][2], (2) a further decrease in the overlap of T-T absorption and fluorescence spectral regions, and (3) an increased photostability. This led to the preparation of pentamethylpyrromethene-BF₂ complex (4,4-difluoro-1,3,5,7,8-pentamethyl-4-bora-3a,4a-diaza-s-indacene) (PMP-BF₂). Some of its laser and other spectroscopic properties were obtained.



Pentamethylpyrromethene-BF₂ Complex
PMP-BF₂



Disodium Pentamethylpyrromethene-
2,6-disulfonate-BF₂ Complex
PMPDS-BF₂

2. EXPERIMENTAL

2.1 The dye laser

We used the same small dye laser which had a dye cell 50 mm long and 2.5 mm diameter [3]. Again, the dye laser was used in the static mode (the dye solution not circulating). The absence of an accumulation of detectable amounts of decomposition products in the stagnant solution was evidence for a high order of the laser dye photostability. To record laser output energies, we used a Scientech model 365 power/energy meter.

2.2 The spectroscopic equipment

We used the same spectroscopic equipment to measure the fluorescence spectrum of PMP-BF₂ as in [3]. Also, the same spectroscopic equipment was used as in [3] to record the T-T absorption (triplet optical density OD_T)



Codes	
Dist	Avail and/or Special
A-1	

spectra, employing the 496.5 nm line from an ion argon cw laser. The conventional singlet-singlet (S-S) absorption spectrum [$\epsilon_{492}(\text{CH}_3\text{OH}) = 85,200$] was recorded with Cary 17 and Varian SuperScan 3 UV/visible spectrophotometers.

To obtain triplet extinction coefficients $\epsilon_T(\lambda)$ from the measured $\text{OD}_T(\lambda)$ values we employed McClure's method [4-6]. Because of the limited solubility of PMP-BF₂ in ethanol, we used 2-methyltetrahydrofuran as a glassy solvent.

2.3. The laser dyes and chemicals

The synthesis of PMP-BF₂ and the disodium salt of 1,3,5,7,8-pentamethylpyrromethene-2,6-disulfonic acid-BF₂ complex (PMPDS-BF₂) will be reported elsewhere. Coumarin 545 was obtained from Exciton and rhodamine 6G from the Eastman Kodak Company. Ethanol was purchased from the U.S. Industrial Company, 2-methyltetrahydrofuran from Lancaster Synthesis, and N,N-dimethylformamide from the Aldrich Chemical Company.

3. RESULTS AND DISCUSSION

3.1 Pentamethylpyrromethene-BF₂ complex

In figure 1 we present the energy output ΔE (in mJ) of a 1.5×10^{-4} molar solution of PMP-BF₂ in ethanol as a function of flashlamp pump energy E (in J). In the same figure, we also show the energy output of a 2×10^{-4} molar solution of coumarin 545 in ethanol and a 2×10^{-4} molar solution of rhodamine 6G also dissolved in ethanol. PMP-BF₂ lases BB at 546 nm, coumarin 545 at 546 nm, and rhodamine 6G at 595 nm. From Figure 1 it is apparent that PMP-BF₂ lases about 300% more efficiently than coumarin 545 and about 20% less efficiently than rhodamine 6G. PMP-BF₂ was about as photostable as rhodamine 6G.

Figure 2 shows the fluorescence, S-S absorption and T-T absorption of PMP-BF₂. A striking feature is the exceptionally low value of the triplet extinction coefficient $\epsilon_T \sim 1 \times 10^3$ L/mole cm over the laser action spectral region.

In a comparison evaluation with acridine orange ($Q_F = 0.46$) fluorescence quantum yields (Q_F) were obtained for PMP-BF₂ (0.73) PMPDS-BF₂ (0.98) and TMP-BF₂ (0.83, compare 0.80 [2]).*

3.2 Disodium Pentamethylpyrromethene-2,6-disulfonate-BF₂-Complex (PMPDS-BF₂)

PMP-BF₂ was partially soluble in ethanol/water mixtures and other aqueous solvents, but had limited solubility in water only. PMPDS-BF₂ was readily soluble in water, but showed poor photostability. It showed laser action BB at 555 nm that was about 50% as efficient as that obtained from PMP-BF₂. The salt dissolved in dimethylformamide showed greater photostability and lased BB at 577 nm with about 50% of the efficiency for PMP-BF₂. In methanol PMPDS-BF₂ (2×10^{-4} M) lased at 551 nm. A comparison of the laser energy output ΔE of ethanol solutions of selected laser dyes as a function of flashlamp pump energy E , figure 1, shows PMPDS-BF₂ to be about 75% as efficient as PMP-BF₂ in ethanol. In methanol PMPDS-BF₂ was vastly superior to rhodamine 6G in photostability.

3.3 Discussion

Our experimental results on the lasing efficiency of PMP-BF₂ and PMPDS-BF₂ places these new laser dyes and TMP-BF₂ among the most impressive laser dyes known. Nevertheless, we are surprised that they did not lase with even greater efficiency. From Figure 2 it follows that PMP-BF₂ is the dye

* We are indebted to Dr. Scott Whittenburg for these Q_F determinations.

with the lowest $\epsilon_T(\lambda_{las})$ of any laser dye known [6]. The prolonged laser action efficiency of these three dyes, TMP-BF₂, PMP-BF₂, and PMPDS-BF₂, can be attributed to a combination of their high extinction coefficients, moderately high fluorescence quantum yields, minimal S-S and T-T absorption in the fluorescence spectral region, and superior photostability in selected solvents. Photodegradation of PMP-BF₂ from flashlamp irradiation was not detected. Continuous irradiation of PMP-BF₂ (10⁻⁴ M) in methanol by a Westinghouse sunlamp (275 w) at a distance of about 20 cm. from the flask brought about the disappearance of fluorescence after one week. In a similar experiment in which methanol as solvent was replaced by methylene chloride fluorescence had not disappeared after one month. To obtain increased prolonged power efficiency tailor made derivatives of the pyrromethene-BF₂ complex that offer further improvement in these characteristic features are sought.

Acknowledgements:

The authors gratefully acknowledge the support from the Office of Naval Research. One of the authors (TGP) wishes to thank the Naval Ocean Systems Independent Research program; another author (JHB) thanks the Board of Regents Louisiana Education Quality Support Fund under contract LEQSF(86-87)-RD-B-06.

REFERENCES

1. T. G. Pavlopoulos, M. Shah, and J. H. Boyer, Appl. Optics, 15 December 1988.
2. E. Vos de Wael, J. A. Pardoën, J. A. van Koeveringe, and J. Lugtenburg, Rec., Trav. Chim. Pays-Bas 96, 306 (1977).
3. T. G. Pavlopoulos, J. H. Boyer, I. R. Politzer, and C. M. Lau, J. Appl Phys. 60, 4028 (1986).
4. D. J. McClure, J. Chem. Phys. 19, 670 (1951).
5. T. G. Pavlopoulos, Spectrochim. Acta, 43A, 1201 (1987).
6. T. G. Pavlopoulos and D. J. Golich, J. Appl. Phys. 64, 521 (1988).

Figure 1.

Laser energy output ΔE (in mJoule) as a function of energy E (in Joule) of a 1.5×10^{-4} molar solution of the pentamethylpyrromethene- BF_2 complex (PMP- BF_2) dissolved in ethanol, a 2×10^{-4} molar solution of Coumarin 545 dissolved in ethanol, a 2×10^{-4} molar solution of Rhodamine 6G dissolved in ethanol, and a 2×10^{-4} molar solution of pentamethylpyrromethene- BF_2 sulfonate (PMP- BF_2 -sulfonate) also dissolved in ethanol.





Figure 2.

Absorption (S-S) and fluorescence Fl spectra of pentamethylpyrromethene- BF_2 complex (PMP- BF_2) dissolved in ethanol. The T-T absorption spectrum was recorded at 77 K, employing a 1×10^{-4} molar solution of 2-methyltetrahydrofuran as a glassy solvent.



TECHNICAL REPORT DISTRIBUTION LIST, GENERAL

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Chemistry Division, Code 1113 800 North Quincy Street Arlington, VA 22217-5000	3	Dr. Ronald L. Atkins Chemistry Division (Code 385) Naval Weapons Center China Lake, CA 93555-6001	1
Commanding Officer Naval Weapons Support Center Attn: Dr. Bernard E. Douda Crane, IN 47522-5050	1	Chief of Naval Research Special Assistant for Marine Corps Matters Code OOMC 800 North Quincy Street Arlington, VA 22217-5000	1
Dr. Richard W. Drisko Naval Civil Engineering Laboratory Code L52 Port Hueneme, California 93043	1	Dr. Bernadette Eichinger Naval Ship Systems Engineering Station Code 053 Philadelphia Naval Base Philadelphia, PA 19112	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	2 <u>high</u> <u>quality</u>		
David Taylor Research Center Dr. Eugene C. Fischer Annapolis, MD 21402-5067	1	Dr. Sachio Yamamoto Naval Ocean Systems Center Code 52 San Diego, CA 92152-5000	1
Dr. James S. Murday Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	David Taylor Research Center Dr. Harold H. Singerman Annapolis, MD 21402-5067 ATTN: Code 283	1